




Contents lists available at ScienceDirect

International Journal of Cardiology

journal homepage: www.elsevier.com/locate/ijcard

Blood transfusion and ischaemic outcomes according to anemia and bleeding in patients with non-ST-segment elevation acute coronary syndromes: Insights from the TAO randomized clinical trial

P. Deharo^{a,b,c}, G. Ducrocq^{d,e,f}, C. Bode^g, M. Cohen^h, T. Cuisset^a, S.R. Mehtaⁱ, C.V. Pollack^j, S.D. Wiviott^k, S.V. Rao^l, J.W. Jukema^{m,p}, A. Erglisⁿ, T. Moccetti^o, Y. Elbez^{d,e,f}, P.G. Steg^{q,r,*}

^a Département de Cardiologie, CHU Timone, Marseille F-13385, France

^b Aix Marseille Univ, Inserm, Inra, C2VN, Marseille, France

^c Aix-Marseille Université, Faculté de Médecine, F-13385 Marseille, France

^d Hôpital Bichat, Assistance Publique-Hôpitaux de Paris, Paris, France

^e Université Paris-Diderot, Département Hospitalo-Universitaire FIRE, Sorbonne-Paris Cité, Paris, France

^f FACT (French Alliance for Cardiovascular clinical Trials), an F-CRIN network, NHLL, Royal Brompton Hospital, Imperial College, London, United Kingdom

^g Heart Center Freiburg University, Cardiology and Angiology I, Faculty of Medicine, Freiburg, Germany

^h Newark Beth Israel Medical Centre, Newark NJ and Rutgers-New Jersey Medical School, Newark, New Jersey, USA

ⁱ McMaster University and the Population Health Research Institute, Hamilton Health Sciences, Hamilton, Ontario, Canada

^j Department of Emergency Medicine, Thomas Jefferson University, Philadelphia, Pennsylvania, USA

^k TIMI Study Group, Division of Cardiovascular Medicine, Brigham and Women's Hospital, Harvard Medical School, Boston, Massachusetts, USA

^l Department of Cardiology, Duke Clinical Research Institute, Durham, NC, USA

^m Department of Cardiology, Leiden University Medical Center, Leiden, The Netherlands

ⁿ University of Latvia, Pauls Stradins Clinical University Hospital, Riga, Latvia

^o Electrophysiology Unit, Department of Cardiology, Fondazione Cardiocentro Ticino, via Tesserete 48, 6900 Lugano, Switzerland

^p The Netherlands Heart Institute, Utrecht, the Netherlands

^q French Alliance for Cardiovascular Trials, Département Hospitalo-Universitaire FIRE, AP-HP, Hôpital Bichat, Université de Paris, Institut National de la Santé et de la Recherche Médicale, Paris, France

^r National Heart and Lung Institute, Imperial College, Royal Brompton Hospital, London, UK

ARTICLE INFO

Article history:

Received 10 December 2019

Received in revised form 1 April 2020

Accepted 12 June 2020

Available online xxxx

Keywords:

Non ST elevation Myocardial Infarction
percutaneous coronary intervention
Blood transfusion

ABSTRACT

Background: The benefits and risks of blood transfusion in patients with acute myocardial infarction who are anemic or who experience bleeding are debated. We sought to study the association between blood transfusion and ischemic outcomes according to haemoglobin nadir and bleeding status in patients with NST-elevation myocardial infarction (NSTEMI).

Methods: The TAO trial randomized patients with NSTEMI and coronary angiogram scheduled within 72h to heparin plus eptifibatid versus otamixaban. After exclusion of patients who underwent coronary artery bypass surgery, patients were categorized according to transfusion status considering transfusion as a time-varying covariate. The primary ischemic outcome was the composite of all-cause death or MI within 180 days of randomization. Subgroup analyses were performed according to pre-transfusion hemoglobin nadir and bleeding status. **Results:** 12,547 patients were enrolled. Among these, blood transfusion was used in 489 (3.9%) patients. Patients who received transfusion had a higher rate of death or MI (29.9% vs. 8.1%, $p < 0.01$). This excess risk persisted after adjustment on GRACE score and nadir of hemoglobin (HR 3.36 95%CI 2.63–4.29 $p < 0.01$). Subgroup analyses showed that blood transfusion was associated with a higher risk in patients without overt bleeding (adjusted HR 6.25 vs. 2.85; p -interaction 0.001) as well as in those with hemoglobin nadir > 9.0 g/dl (HR 4.01; p -interaction < 0.0001).

Conclusion: In patients with NSTEMI, blood transfusion was associated with an overall increased risk of ischaemic events. However, this was mainly driven by patients without overt bleeding and those hemoglobin nadir > 9.0 g/dl. This suggests possible harm of transfusion in those groups.

1. Introduction

In patients with acute coronary syndrome (ACS), the presence of anemia, with or without active bleeding, may jeopardize oxygen

* Corresponding author at: Hôpital Bichat, Department of cardiology, 46 rue Henri Huchard, 75018 Paris, France.

E-mail address: gabriel.steg@aphp.fr (P.G. Steg).

delivery to the myocardium, [1–3]. Red blood cell (RBC) transfusion is the only treatment available that rapidly increases hemoglobin level [4]. However, RBC transfusions are also associated with potentially deleterious effects in the acute phase of an ACS [5–7]. Therefore, the indications of RBC transfusion are still debated and clinical management of anemia or bleeding varies widely in this population [8].

The most recent guidelines from the American College of Cardiology/American Heart Association and of the European Society of cardiology for the management of patients with NSTEMI both recommend a restrictive use of transfusion with avoidance of routine RBC transfusion in patients with hemoglobin levels over 7.0 g/dL for European (Class IIb Level C) and 8.0 g/dl (Class III Level B for transfusion if hemoglobin > 8.0 g/dl) for American [9,10]. However, those guidelines are not based on randomized data but on observational studies of outcomes in patients requiring transfusion in the setting of myocardial infarction (MI). Yet, these studies have yielded inconsistent results, with transfusion being associated with either improved or worsened subsequent cardiovascular outcomes [11,12].

We hypothesized that the presence or absence of active bleeding and the severity of anemia may affect the benefits and risks of RBC transfusion in patients with ACS.

Using a large contemporary trial cohort, derived from a recent international randomized trial in NSTEMI patients, we aimed to assess the association between blood transfusion and ischaemic events overall as well as according to hemoglobin nadir and bleeding status.

2. Methods

2.1. Study population

The methods and results of the TAO trial (Treatment of Acute Coronary Syndrome With Otamixaban) have been previously described. Briefly, TAO was a large international trial randomizing patients with moderate to high-risk NSTEMI with coronary angiography planned in the first 72 hours, to heparin plus eptifibatide versus otamixaban. Patients were randomized to the UFH plus eptifibatide group or to 1 of 2 otamixaban dosing groups (intravenous bolus of 0.080 mg/kg followed by an infusion of either 0.100mg/kg per hour or 0.140mg/kg per hour) in a 1:1:1 ratio. A planned interim analysis performed after 1969 randomized patients in each group led the investigators to define the higher-dose otamixaban as optimal, discontinue enrolment in the lower dose and to continue enrolment in the higher dose until study end in a 1:1 ratio compared to placebo.

Eligibility criteria for the study were patients with NSTEMI scheduled to undergo an invasive strategy (angiography and PCI, if indicated, to be performed at the latest within 72 hours of randomization).

The main exclusion criteria were a revascularization procedure already performed for the qualifying event; acute ST-segment elevation myocardial infarction; receipt of a therapeutic dose of injectable anticoagulant for more than 24 hours before randomization; or treatment with abciximab. The main results of the trial have been previously published: otamixaban did not reduce the rate of the primary outcomes of death plus MI but did increase bleeding in comparison with heparin plus eptifibatide [13,14].

PCI is currently the main revascularisation modality in patients with NSTEMI [15–17]. The use of CABG is currently marginal in this setting and is associated with higher bleeding rate [15–18]. In patients undergoing CABG, the use of transfusion is much more frequent than in the overall NSTEMI population (between 30 to 90% according to local protocols) and has recently been evaluated in a dedicated randomized trial [18]. For those reasons, patients who underwent CABG as treatment of the index event were excluded from the present analysis.

2.2. Bleeding definition and transfusion

Bleeding events were reported if clinically overt, of unanticipated/unexpected quantity, prompting medical attention, evaluation or

treatment. Bleeds were categorized using the Thrombosis in Myocardial Infarction (TIMI) classification into major and minor bleeding [19]. In case of reported bleeding, hemoglobin and hematocrit before and after the event were reported [13]. Hemoglobin nadir is defined as the lowest hemoglobin value and was categorized as ≤ 7.0 , 7.1–8.0, 8.1–9.0, 9.1–10.0, >10.0 g/dl.

The indication for transfusion and the number of packed red blood cells or whole blood units transfused were left to the discretion of the treating physicians. The number of red blood cell units transfused was reported.

2.3. Outcomes

The primary ischemic outcome studied was the same as the primary endpoint of the TAO trial: a composite of all-cause death or new MI. For the present analysis, the longest follow-up available was considered for the primary endpoint (i.e. 180 days compared to 7 days for the TAO trial).

Secondary efficacy outcome measures included the same composite ischaemic endpoint at day 30. All-cause death, cardiovascular death, non-fatal MI, stroke, rehospitalization or prolongation of the hospitalization due to MI and stent thrombosis (ST) were analysed separately at day 30. ST were categorized according to the Academic Research Consortium classification and MIs were categorized according to the 2007 universal definition [20,21].

Efficacy outcomes, were adjudicated by a central clinical event committee (TIMI Study Group); the committee members were unaware of study treatment assignments.

In order to explore potential mechanisms in outcome differences, we evaluated the incidence of antiplatelet cessation according to transfusion status, bleeding and hemoglobin nadir. All antiplatelet agents were considered (aspirin, clopidogrel and ticagrelor). A patient fulfilled criteria for antiplatelet cessation when receiving a given antiplatelet at the time of randomization but not at any subsequent time point of index hospitalization.

2.4. Ethics

All patients provided written informed consent. In each country, the study was approved by ethics committees in accordance with local guidelines.

2.5. Statistical analysis

Patients were categorized into two groups according to their transfusion status during entire study follow up. Descriptive statistics are reported as means \pm standard deviations for continuous variables and patient numbers with percentages for categorical variables.

Categorical variables across groups were compared by chi-square tests and continuous variables by analyses of variance. Events rates comparisons across groups were performed by chi-square tests. Primary and secondary outcomes were adjusted according to several multivariable logistic models including predefined variables and considering transfusion as a time-varying covariate. Model 1 was adjusted on GRACE score. Model 2 on GRACE score and nadir of hemoglobin. Model 3 included adjustment on GRACE score, region, sex, atrial fibrillation, peripheral artery disease and smoking status. Outcomes were adjusted on models 1, 2 and 3. We also present in supplemental analysis an analysis of the primary endpoint and according to bleeding status and nadir of hemoglobin with addition to the model 3 (GRACE score, region, sex, atrial fibrillation, peripheral artery disease, smoking status) of body mass index, treatment with percutaneous coronary intervention and TAO trial allocated treatment (i.e. heparin and eptifibatide vs. otamixaban).

Given the lower number of events, subgroup analyses according to bleeding status were adjusted on model 1 only. For the subgroup

analysis according to nadir hemoglobin, an additional model (model 4) was built for adjustment on GRACE score and baseline hemoglobin.

Comparisons between transfused and non-transfused patients were performed according to bleeding status (TIMI minor and major or no bleeding) occurring within 7 days after randomization, and according to the hemoglobin nadir.

The interactions between bleeding subgroups, as well as hemoglobin nadir subgroups and transfusion status were assessed by introducing interaction terms in the logistic models.

P values were reported for all statistical tests, with a cut off of 0.05 to consider a statistical significance, except for p values of interaction where the cut off of significance was set as 0.10. All statistical analyses were performed using the statistical software SAS, version 9.3 (Statistical Analyses System, SAS Institute Inc., Cary, NC, USA).

3. Results

3.1. Baseline and procedural characteristics

In the TAO trial, 13,229 patients with NSTEMI were randomized. Of these, 682 (5.1%) underwent CABG and were excluded from analysis. Within the study population of 12,547 patients, 12,058 (96.1%) did not receive any transfusion during the study follow up, while 489 patients (3.9%) received at least one RBC transfusion (Supplemental Figure 1).

Table 1 describes the baseline characteristics of the study population. Patients with transfusion were frailer (older and with lower body mass index) and sicker (higher Killip and GRACE scores and lower creatinine clearance). Vascular disease, cardiovascular risk factors and atrial fibrillation were more prevalent in this group. These patients also received less frequently aspirin or P2Y12 blockers and had lower rates of PCI following the acute MI ($p=0.05$). Finally, cessation of an antiplatelet agent was reported in 20.3% ($n=97$) of transfused patients versus 5.3% ($n=627$) of non-transfused patients ($p<0.001$) (Supplemental table 1).

3.2. Outcomes

3.2.1. Ischaemic outcomes

Unadjusted outcomes are presented in Supplemental Table 2. In the overall population, at 180 days, the primary efficacy endpoint occurred in 719 patients (6.0%) in the non-transfused cohort and 116 (23.7%) in the transfused cohort ($p<0.001$).

When transfusion status was introduced as a time-varying covariate in the Cox model the unadjusted HR for the primary endpoint was 7.35 (95%CI 6.00-9.015, $p<0.01$) (Fig. 1).

The increase in ischaemic risk associated with transfusion persisted after adjustment across all models: model 1 (GRACE score), model 2 (GRACE score and nadir of haemoglobin) as well on model 3 (GRACE score, region, sex, atrial fibrillation, peripheral artery disease history, smoking status) (adjusted HR 6.17 (4.98-7.65), $p<0.001$; 3.36 (2.63-4.29), $p<0.001$ and 5.65 (4.54-7.03), $p<0.001$ respectively) (Fig. 1). In each model, the rates of MI and of death were higher in patients who had received RBC transfusion than in patients who were not transfused (Supplemental figure 2).

3.3. Ischaemic outcomes according to bleeding status

In the transfusion group, 85 patients (17.4%) experienced a TIMI major (65 patients) or minor (22 patients) bleed during follow up, versus 231 (1.9%) (19 TIMI major and 214 TIMI minor) in the non-transfused group.

Unadjusted outcomes according to bleeding status are presented in Figure 2. Patients transfused despite not having experienced a bleeding event had a higher risk of the primary endpoint than those with bleeding (HR 7.49 95%CI (5.97-9.41) vs. HR 3.20 95%CI (1.93-5.32)). After

adjustment on model 1 (GRACE score), this excess risk persisted (HR 6.25 (4.91-7.95) vs HR 2.85 (1.70-4.80) p value for interaction 0.0066) (Fig. 2). This increased incidence of the primary endpoint was driven by an increased risk of recurrent MI in patients with transfusion (5.52 (4.05-7.53) vs 2.24 (1.11-4.55) p value for interaction 0.02) whereas there was no excess mortality risk (5.37 (3.80-7.60) vs 6.56 (3.35-12.82) p value for interaction 0.549) (Supplemental figure 3).

Within the transfused cohort, no difference in any antiplatelet cessation according to bleeding status was reported: 16 (19.3%) patients with bleeding vs. 81 (20.5%) patients without bleeding ($p=0.80$).

3.3.1. Ischaemic outcomes according to hemoglobin nadir

To further explore the association between RBC transfusion and risk of death and MI we stratified analyses according to the nadir of haemoglobin.

Unadjusted outcomes are presented in Fig. 3 and show worse ischaemic outcomes for patients who received transfusion but had the highest haemoglobin nadir (p interaction 0.0018).

After adjustment on GRACE score (model 1) and GRACE score and baseline hemoglobin (model 4), a graded association between blood transfusion and the primary endpoint was observed for patients with nadir of haemoglobin > 9.0 g/dl (p value for interaction 0.006 in model 1 and <0.0001 in model 2). Above a certain Hb value (Hb > 8.0 g/dl in model 1 and haemoglobin > 9 g/dl in model 2) RBC transfusion was associated with a higher risk of all cause death or MI (Fig. 3). The two components of the primary endpoint are presented in supplemental figures 4 and 5. In model 1, the rates of recurrent non-fatal MI were significantly influenced by RBC transfusion (p value interaction 0.0009) while in model 2 both all-cause death and MI were independently associated with RBC transfusion (p 0.008 and $p<0.001$ respectively).

Within the transfused cohort, hemoglobin nadir was not associated with antiplatelet cessation ($p=0.09$).

Supplemental figure 6 presents the results adjusted on GRACE score, region, sex, AF, PAD history, smoking status body mass index, treatment with percutaneous coronary intervention and TAO trial allocated treatment (i.e. heparin and eptifibatid vs. otamixaban) showing similar results than in the main analysis.

4. Discussion

In the present analysis, RBC transfusion in NSTEMI patients was associated with increased risks of death and MI. This increase in ischemic events was mainly driven by patients transfused without overt bleeding and those with hemoglobin nadir > 9.0 g/dl.

In theory, RBC transfusion, because it quickly increases hemoglobin level, should increase oxygen delivery to ischemic myocardium and limit myocardial injury during the acute phase of MI. However, data suggest that RBCs have high oxygen affinity and low 2,3-diphosphoglycerate, and, as a consequence, oxygen delivery may in fact not be improved in patients receiving RBC transfusions [22,23]. In addition, during storage, packed RBCs are rapidly depleted of nitric oxide, leading to attenuation of physiologic vasodilation in hypoxic areas, impairment of erythrocyte function and deformability, and disruption of normal oxygen delivery in the microcirculation [24-26]. There is also evidence of increased platelet reactivity resulting from RBC transfusion [27]. Therefore, transfusion may actually be deleterious at least in some patients during acute coronary syndrome. There is a need to better understand whether RBC transfusion is associated with benefit or harm and in which types of patients.

Currently, both European and American guidelines recommend a restrictive approach to transfusion in patients with ACS which implies withholding transfusion if the hemoglobin level exceeds 7.0 g/dL to 8.0 g/dL [9,10]. However, few clinical data are available to support this recommendation. The only two randomized studies performed in this setting were underpowered (45 and 110 patients respectively) and showed conflicting results [28,29]. Currently available randomized

Table 1
Baseline characteristics and procedural characteristics of the population, according to transfusion status.

	No transfusion N= 12058	Transfusion N= 489	p-value
Clinical presentation:			
Age, mean (sd) y	61.66 (11.22)	66.22 (11.34)	<0.001
Body weight, mean (sd) kg	80.75 (16.7)	75.92 (15.7)	<0.001
Body mass index, mean (sd), kg/m ²	28.26 (5.02)	27.7 (4.92)	0.007
Male gender, n (%)	8430 (69.9%)	279 (57.1%)	<0.001
Systolic BP, mean (SD), mmHg	131.88 (19.55)	132.13 (21.91)	0.941
Diastolic BP, mean (SD), mmHg	77.09 (11.95)	75.1 (13.53)	0.002
Heart rate, mean (SD), bpm	72.01 (12.92)	76.08 (14.5)	<0.001
Killip class, n (%):			<0.001
I	11218 (93.5%)	405 (82.8%)	
II	675 (5.6%)	63 (12.9%)	
III	98 (0.8%)	19 (3.9%)	
IV	10 (0.1%)	2 (0.4%)	
Cardiac arrest at admission, n (%)	70 (0.6%)	3 (0.6%)	0.930
Atrial fibrillation on first ECG, n (%)	268 (2.2%)	20 (4.1%)	<0.001
Region, n (%):			<0.001
North America	1455 (12.1%)	84 (17.2%)	
Western Europe	2365 (19.6%)	75 (15.3%)	
Eastern Europe	4185 (34.7%)	107 (21.9%)	
Asia	1054 (8.7%)	49 (10%)	
Other	2999 (24.9%)	174 (35.6%)	
Medical History, n (%):			
Coronary artery disease	4856 (40.3%)	238 (48.7%)	<0.001
Myocardial infarction	2324 (19.3%)	108 (22.1%)	0.124
Coronary artery bypass graft (CABG)	802 (6.7%)	48 (9.8%)	0.006
Percutaneous coronary intervention (PCI)	442 (3.7%)	19 (3.9%)	0.799
Stroke/TIA	617 (5.1%)	44 (9%)	<0.001
Peripheral arterial disease	556 (4.6%)	44 (9%)	<0.001
Carotid endarterectomy/stenting	113 (0.9%)	9 (1.8%)	0.045
Congestive heart failure	609 (5.1%)	57 (11.7%)	<0.001
Hypertension	8499 (70.5%)	408 (83.4%)	<0.001
Hypercholesterolemia	6392 (53%)	271 (55.4%)	0.295
Diabetes mellitus	3315 (27.5%)	194 (39.7%)	<0.001
Current smoker	4098 (34%)	121 (24.7%)	<0.001
Family history of coronary artery disease (CAD)	3882 (32.2%)	153 (31.3%)	0.674
Alcohol habits: n (%):			<0.001
Never	6789 (56.4%)	337 (68.9%)	
At least monthly	2153 (17.9%)	67 (13.7%)	
At least weekly	1976 (16.4%)	50 (10.2%)	
At least daily	35 (7.2%)	35 (7.2%)	
Inclusion criteria, n (%):			
Biomarker elevation	10608 (88%)	434 (88.8%)	0.603
ECG changes	4836 (40.1%)	240 (49.1%)	0.003
GRACE risk score at baseline, n (%):			<0.001
<96	1675 (14.9%)	37 (7.9%)	
96 - 112	1989 (17.7%)	47 (10%)	
113 - 133	2996 (26.7%)	100 (21.2%)	
>133	4556 (40.6%)	287 (60.9%)	
TIMI risk score at baseline, n (%):			<0.001
0-2	3845 (31.9%)	97 (19.8%)	
3-4	5754 (47.7%)	258 (52.8%)	
5-7	2459 (20.4%)	134 (27.4%)	
Creatinine clearance, median (std), mL/min:	94.85 (36.61)	77.06 (35.74)	<0.0001
Anticoagulation during the PCI, n (%):			0.012
Unfractionated Heparin + eptifibatide	5001 (41.5%)	170 (34.8%)	
Otamixaban 0.100	2414 (20%)	107 (21.9%)	
Otamixaban 0.140	4643 (38.5%)	212 (43.4%)	
Antiplatelet therapy received between randomization and discharge, n (%):			
Aspirin	11834 (98.1%)	462 (94.5%)	<0.001
Oral ADP receptor antagonist:			
Clopidogrel	10559 (87.6%)	371 (75.9%)	<0.001
Prasugrel	608 (5%)	10 (2%)	0.002
Ticagrelor	416 (3.4%)	14 (2.9%)	0.484
Other medications between randomization and discharge, n (%):			
Statins	11187 (92.8%)	450 (92%)	0.529
ACEI	9453 (78.4%)	367 (75.1%)	0.078
betablockers	9997 (82.9%)	426 (87.1%)	0.014
Management during the index admission, n (%):			
Percutaneous coronary intervention (PCI)	8474 (70.3%)	182 (37.2%)	<0.001
Neither PCI nor CABG	3584 (29.7%)	307 (62.8%)	<0.001
Duration of the index hospitalization, mean (sd) days	5.38 (3.38)	9.16 (7.88)	<0.001

GRACE: Global Registry of Acute Coronary Events; TIMI: Thrombolysis in Myocardial Infarction;
ECG: electrocardiogram; ACEI: angiotensin-converting-enzyme inhibitor




Fig. 1. Unadjusted and adjusted hazard ratio of primary outcome at 180 days comparing transfused vs. not transfused patients (reference). Adjustment covariates are: GRACE score (model 1) GRACE score and nadir of hemoglobin (model 2) and GRACE score, Region, Sex, AF, PAD history, Smoking status (model 3). Transfusion status has been introduced as time varying covariate in the model.

data on blood transfusion in MI patients are therefore scarce and rely on very small datasets, precluding any definitive conclusion.

A recent meta-analysis of randomized trials examining optimal thresholds for RBC transfusion among inpatients with various conditions found no difference in 30 day mortality between liberal and restrictive transfusion strategies [30]. However, the specific sub-analysis of the 2 small randomized trials in ACS patients showed a trend favouring liberal strategy (RR 3.88; 95% CI 0.83–18.13). This subanalysis only included 154 patients and is underpowered as attested by the large confidence interval. However, those data suggests that ACS might be a specific setting in which the balance of benefit and risk of transfusion may differ from other clinical situations and deserves to be analyzed separately.

Most of the available clinical evidence derives from observational data, which tend to support a restrictive rather than a liberal transfusion strategy in ACS [11,12,31,32, and]. Three large meta-analyses of observational data, including both STEMI and NSTEMI patients, have concluded that RBC transfusion is associated with higher short and long-term mortality after ACS [11,12,33, and]. In 2 out of these 3 meta-analyses, RBC transfusion at hemoglobin below 8 g/dL appeared to be beneficial, and its use at hemoglobin above 10.0 g/dL was associated with an increased risk of mortality [11,33]. One of the main hypotheses for such conflicting results is that both anemia and bleeding, which drive transfusion

indications, are themselves associated with a worse prognosis in patients with MI, and confound the association of transfusion with outcomes [34]. There might be some heterogeneity in the effect of transfusion in ACS patients according to different clinical characteristics. Recent observational data suggest worse outcome in patients with RBC transfusion only in case of STEMI but no difference in case of NSTEMI [8].

To the best of our knowledge, a differential effect of transfusion according to the presence of active bleeding has never been explored. In the present analysis, we observed that, in NSTEMI patients, RBC transfusion was associated with increased ischemic hazard mostly in patients without bleeding or with high hemoglobin levels. After statistical adjustment, we identified significant interaction between transfusion and bleeding, meaning that patients transfused without having experienced significant bleeding had a worse prognosis. Same results were observed with nadir of hemoglobin above 9.0 g/dl.

It is conceivable that in patient with active bleeding and lowest hemoglobin levels the balance between beneficial and deleterious effects of RBC transfusion is positive, whereas it is negative in others. A confounding component of the increased ischemic risk in the transfused group could be the need to discontinue antithrombotic drugs when bleeding or anemia occur. An increase in platelet reactivity combined with discontinuation in dual antiplatelet therapy could explain the higher risk for recurrent MI after transfusion [26,27]. In the present




Fig. 2. Unadjusted and adjusted hazard ratio of primary endpoint at 180 days comparing transfused vs. not transfused patients stratified by status of bleeding events occurring in the 7 days after randomization. Transfusion status has been introduced as time varying covariate in the model. The analysis was performed adjusted with GRACE score (model 1).




Fig. 3. Unadjusted and adjusted hazard ratio of primary endpoint at 180 days comparing transfused vs. not transfused patients stratified by group of nadir of hemoglobin. Transfusion status has been introduced as time varying covariate in the model. The primary efficacy outcome, all cause death and non-fatal MI was analyzed until 180 days. The analysis was performed adjusted with GRACE score (model 1) GRACE score and Baseline Hemoglobin (model 4)

analysis, we observed a nearly four-time higher rate in antiplatelet cessation in patients receiving transfusion compared to non-transfused patients. This can at least partly explain the increased ischemic risk in transfused patients.

We observed, however, no differences in antiplatelet cessation in transfused patients according to bleeding or hemoglobin status. This suggests that the heterogeneity of outcomes observed in these subgroups is more likely related to the effect of transfusion than to the confounding component of antiplatelet cessation.

Patients undergoing surgical revascularization were excluded from this analysis. The question of transfusion in patients undergoing CABG has been specifically addressed in several adequately powered randomized trial and recent evidence is in favor of a restrictive RBC transfusion approach in this setting [18]. In our cohort we observed that in the transfused cohort the majority of patients were treated medically. While, in the non-transfused cohort NSTEMI patients were more often treated invasively. Therefore, the occurrence of transfusion do impact the treatment strategy in acute coronary syndrome. Indeed, patients undergoing transfusion are more often managed medically.

4.1.1. Limitations

Although multivariable adjustment was used to correct for measured differences between the groups, we cannot exclude the presence of unmeasured confounders. It is important to note that this study is a post hoc analysis, and our findings should therefore be interpreted as hypothesis-generating only. The indications for transfusion were not prespecified in TAO trial and varied across centers.

Patients with RBC transfusion were at higher baseline risk and had different antithrombotic regimen characterized by reduced use of aspirin and clopidogrel.

Consequently, only large randomized controlled trials will be able to address this issue definitively. Currently, 2 randomized trials are ongoing and will provide high quality evidence regarding optimal strategies for RBC transfusion in ACS patients. The REALITY (REstrictive And

Liberal Transfusion Strategies in Patients With Acute myocardial Infarction) trial (NCT02648113) is randomizing 630 ACS patients in a restrictive transfusion strategy (RBC transfusion if hemoglobin < 8.0 g/dl with objective 8.0 to 10.0 g/dl) vs. liberal strategy (RBC transfusion if hemoglobin < 10.0 g/dl with objective of 11.0 g/dl) [35] and will evaluate a 30-day composite of all-cause death, non-fatal stroke, nonfatal recurrent MI, and emergency revascularization. The MINT (Myocardial Ischemia and Transfusion) trial (NCT02981407) aims to randomize 3500 patients with MI and anemia to a restrictive (hemoglobin < 8.0 g/dl) vs. liberal strategy (hemoglobin < 10.0 g/dl) [36]. The primary outcome will be a 30-day composite of all-cause mortality or nonfatal MI.

5. Conclusion

In patients with NSTEMI, RBC transfusion was associated with an increase of risk of death or MI at 180 days. This worse outcome appeared more pronounced in case of transfusion without overt bleeding and for hemoglobin level > 9.0 g/dl. This suggests that it may be prudent to refrain from RBC transfusion in those patients who do not have overt bleeding and have preserved hemoglobin levels. Randomized trials are required to better define patients who will benefit from RBC transfusion after a NSTEMI.

Acknowledgements

To our nurses' team and technicians in executing this study. To the TAO investigators.

Disclosures

PG Steg Research grants: Amarin, Bayer, Sanofi, and Servier Clinical trials (Steering committee or CEC or DMC), speaker or consultant : Amarin, AstraZeneca, Bayer, Boehringer Ingelheim, Bristol-Myers Squibb, Idorsia, Mylan, Novo-Nordisk, Novartis, Pfizer, Sanofi, Servier. CV Pollack Consultant for Sanofi.

J W Jukema : JW Jukema/his department has received research grants from and/or was speaker (with or without lecture fees) on a.o. (CME accredited) meetings sponsored by Amgen, Athera, Astra-Zeneca, Biotronik, Boston Scientific, Dalcor, Daiichi Sankyo, Lilly, Medtronic, Merck-Schering-Plough, Pfizer, Roche, Sanofi Aventis, The Medicine Company, the Netherlands Heart Foundation, CardioVascular Research the Netherlands (CVON), the Netherlands Heart Institute and the European Community Framework KP7 Programme.

M Cohen : speaker Sanofi

SD Wiviott: Sanofi Aventis, AstraZeneca, Merck, Eli Lilly/Daiichi Sankyo, Boehringer Ingelheim.

SR Mehta: Research grants AstraZeneca and Boston scientific;

M Sabatine : to be suppressed as not any more an author for a long time

Sunil V Rao A.Erglis , C Bode, T Cuisset, T Moccetti, G Ducrocq Y Elbez None.

Fundings

TAO trial was funded by Sanofi. For this analysis, no funding source to declare.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.ijcard.2020.06.020>.

References

- [1] M.S. Sabatine, D.A. Morrow, R.P. Giugliano, et al., Association of hemoglobin levels with clinical outcomes in acute coronary syndromes, *Circulation* 26 (2005) 111: 2042-9.
- [2] M. Moscucci, K.A. Fox, C.P. Cannon, et al., Predictors of major bleeding in acute coronary syndromes: the Global Registry of Acute Coronary Events (GRACE), *Euro. Heart J.* 24 (2003) 1815-1823.
- [3] J.W. Eikelboom, S.R. Mehta, S.S. Anand, C. Xie, K.A. Fox, S. Yusuf, Adverse impact of bleeding on prognosis in patients with acute coronary syndromes, *Circulation* 114 (2006) 774-782.
- [4] K.P. Alexander, A.Y. Chen, T.Y. Wang, CRUSADE Investigators, et al., Transfusion practice and outcomes in non-ST-segment elevation acute coronary syndromes, *Am. Heart J.* 155 (2008) 1047-1053.
- [5] X. Yang, K.P. Alexander, A.Y. Chen, CRUSADE Investigators, et al., The implications of blood transfusions for patients with non-ST-segment elevation acute coronary syndromes: results from the CRUSADE National Quality Improvement Initiative, *J. Am. Coll. Cardiol.* 46 (2005) 1490-1495.
- [6] S.V. Rao, J.A. Eikelboom, C.B. Granger, R.A. Harrington, R.M. Califf, J.P. Bassand, Bleeding and blood transfusion issues in patients with non-ST-segment elevation acute coronary syndromes, *Eur. Heart J.* 28 (2007) 1193-1204.
- [7] S.V. Rao, J.G. Jollis, R.A. Harrington, et al., Relationship of blood transfusion and clinical outcomes in patients with acute coronary syndromes, *JAMA* 292 (2004) 1555-1562.
- [8] S. Gili, F. D'Ascenzo, M.F. Lococo, et al., Impact of blood transfusion on in-hospital myocardial infarctions according to patterns of acute coronary syndrome: Insights from the BleeMACS registry, *Int. J. Cardiol.* 221 (2016) 364-370.
- [9] M. Roffi, C. Patrono, J.P. Collet, ESC Scientific Document Group, et al., 2015 ESC Guidelines for the management of acute coronary syndromes in patients presenting without persistent ST-segment elevation: Task Force for the Management of Acute Coronary Syndromes in Patients Presenting without Persistent ST-Segment Elevation of the European Society of Cardiology (ESC), *Eur. Heart J.* 37 (2016) 267-315.
- [10] E.A. Amsterdam, N.K. Wenger, R.G. Brindis, et al., AHA/ACC guideline for the management of patients with Non-ST-elevation acute coronary syndromes: a report of the American College of Cardiology/American Heart Association Task Force on practice guidelines, *J. Am. Coll. Cardiol.* 64 (2014) e139-e228.
- [11] M. Garfinkle, P.R. Lawler, K.B. Filion, M.J. Eisenberg, Red blood cell transfusion and mortality among patients hospitalized for acute coronary syndromes: a systematic review, *Int. J. Cardiol.* 164 (2013) 151-157.
- [12] S. Chatterjee, J. Wetterslev, A. Sharma, E. Lichstein, D. Mukherjee, Association of blood transfusion with increased mortality in myocardial infarction: a meta-analysis and diversity-adjusted study sequential analysis, *JAMA Intern. Med.* 173 (2013) 132-139.
- [13] P.G. Steg, S.R. Mehta, C.V. Pollack Jr., et al., Design and rationale of the treatment of acute coronary syndromes with otamixaban trial: a double-blind triple-dummy 2-stage randomized trial comparing otamixaban to unfractionated heparin and eptifibatid in non-ST-segment elevation acute coronary syndromes with a planned early invasive strategy, *Am. Heart J.* 164 (2012) 817-824 e13.
- [14] P.G. Steg, S.R. Mehta, C.V. Pollack Jr., TAO Investigators, et al., Anticoagulation with otamixaban and ischemic events in non-ST-segment elevation acute coronary syndromes: the TAO randomized clinical trial, *JAMA* 310 (2013) 1145-1155.
- [15] F.J. Neumann, M. Sousa-Uva, A. Ahlsson, et al., ESC/EACTS guidelines on myocardial revascularization. 2018 ESC/EACTS guidelines on myocardial revascularization, *Eur. Heart J.* 40 (2018) 87-165 2019.
- [16] I. Ranasinghe, B. Alprandi-Costa, V. Chow, et al., Risk stratification in the setting of non-ST elevation acute coronary syndromes 1999-2007, *Am. J. Cardiol.* 108 (2011) 617-624.
- [17] T. Fukui, M. Tabata, S. Morita, S. Takahashi, Early and long-term outcomes of coronary artery bypass grafting in patients with acute coronary syndrome versus stable angina pectoris, *J. Thorac. Cardiovasc. Surg.* 145 (2013) 1577-1583.
- [18] C.D. Mazer, R.P. Whitlock, D.A. Fergusson, T.R.I.C.S. Investigators, Perioperative Anesthesia Clinical Trials Group, et al., Six-month outcomes after restrictive or liberal transfusion for cardiac surgery, *N. Engl. J. Med.* 379 (2018) 1224-1233.
- [19] R. Mehran, S.V. Rao, D.L. Bhatt, et al., Standardized bleeding definitions for cardiovascular clinical trials: a consensus report from the Bleeding Academic Research Consortium, *Circulation* 123 (2011) 2736-2747.
- [20] D.E. Cutlip, S. Windecker, R. Mehran, Academic Research Consortium, et al., Clinical end points in coronary stent trials: a case for standardized definitions, *Circulation* 115 (2007) 2344-2351.
- [21] K. Thygesen, J.S. Alpert, H.D. White, Joint ESC/ACC/AHA/WHF task force for the re-definition of myocardial infarction. Universal definition of myocardial infarction, *J. Am. Coll. Cardiol.* 50 (2007) 2173-2195.
- [22] H.G. Welch, K.R. Meenan, L.T. Goodnough, Prudent strategies for elective red blood cell transfusion, *Ann. Intern. Med.* 116 (1992) 393-402.
- [23] M. Casutt, B. Seifert, T. Pasch, E.R. Schmid, M.I. Turina, D.R. Spahn, Factors influencing the individual effects of blood transfusions on oxygen delivery and oxygen consumption, *Crit Care Med.* 27 (1999) 2194-2200.
- [24] S.V. Rao, R.M. Califf, Is old blood bad blood? *Am. Heart J.* 159 (2010) 710-712.
- [25] G.M. Hare, J. Freedman, Mazer C. David, Review article: risks of anemia and related management strategies: can perioperative blood management improve patient safety? *Can. J. Anaesth* 60 (2013) 168-175.
- [26] K.M. Twomley, S.V. Rao, R.C. Becker, Proinflammatory, immunomodulating, and prothrombotic properties of anemia and red blood cell transfusions, *J. Thromb. Thrombolysis* 21 (2006) 167-174.
- [27] J. Silvain, J. Abtan, M. Kerneis, et al., Impact of red blood cell transfusion on platelet aggregation and inflammatory response in anemic coronary and noncoronary patients: the TRANSFUSION-2 study (impact of transfusion of red blood cell on platelet activation and aggregation studied with flow cytometry use and light transmission aggregometry), *J. Am. Coll. Cardiol.* 63 (2014) 1289-1296.
- [28] H.A. Cooper, S.V. Rao, M.D. Greenberg, et al., Conservative versus liberal red cell transfusion in acute myocardial infarction (the CRIT Randomized Pilot Study), *Am. J. Cardiol.* 108 (2011) 1108-1111.
- [29] J.L. Carson, M.M. Brooks, J.D. Abbott, et al., Liberal versus restrictive transfusion thresholds for patients with symptomatic coronary artery disease, *Am. Heart J.* 165 (2013) 964-971 e1.
- [30] J.L. Carson, S.J. Stanworth, J.H. Alexander, et al., Clinical trials evaluating red blood cell transfusion thresholds: an updated systematic review and with additional focus on patients with cardiovascular disease, *Am. Heart J.* 200 (2018) 96-101.
- [31] J.L. Carson, P.A. Carless, P.C. Hébert, Transfusion thresholds and other strategies for guiding allogeneic red blood cell transfusion, *Cochrane Database Syst. Rev.* 2016 (2012) CD002042.
- [32] J.L. Carson, P.A. Carless, P.C. Hébert, Outcomes using lower vs higher hemoglobin thresholds for red blood cell transfusion, *JAMA* 309 (2013) 83-84.
- [33] Y. Wang, X. Shi, R. Du, Y. Chen, Q. Zhang, Impact of red blood cell transfusion on acute coronary syndrome: a meta-analysis, *Intern. Emerg. Med.* 13 (2018) 231-241.
- [34] G. Ducrocq, E. Puymirat, P.G. Steg, et al., Blood transfusion, bleeding, anemia, and survival in patients with acute myocardial infarction: FAST-MI registry, *Am. Heart J.* 170 (2015) 726-734 e2.
- [35] Steg PG. [Clinicaltrials.gov](https://clinicaltrials.gov): cost-effectiveness and cost-utility of liberal vs restrictive red blood cell transfusion strategies in patients with acute myocardial infarction and anaemia (REALITY). NCT02648113, 2017. 2017.
- [36] J.L. Carson, [Clinicaltrials.gov](https://clinicaltrials.gov): myocardial ischemia and transfusion (MINT), NCT02981407 (2017) 2017.